

Effect of Pre-Eclampsia with risk factors for birth weight and neonatal asphyxia: A retrospective study in pregnant women

Awatiful Azza, Esti Yunitasari, Cipto Susilo, Mira Triharini

Faculty of Health Science of Universitas Muhammadiyah Jember, Universitas Airlangga, Mulyorejo, Surabaya City, East Java

Objective: To analyze the risk factors for preeclampsia with birth weight and the incidence of asphyxia.

Methodology: This study used a case-control (retrospective) design with data sources of medical records for 255 mothers who gave birth with a diagnosis of preeclampsia at a referral hospital in East Java, Indonesia the period 2019 to 2020. The indicator for assessing birth weight is seen if it is less than 2500 grams, it is said to be low birth weight, and if it is more than 2500 – 3000 normal body weight. Assessment of neonatal asphyxia uses the Apgar score if the score is 1 – 3 severe asphyxia, 4 – 6 moderate asphyxia, and 7 – 10 mild asphyxia. Research data analysis was carried out using SPSS 21.

Results: The findings of the study on the relationship

between preeclampsia status and newborn body weight obtained a p-value of 0.00, (OR 84.00 95%, CI 27.91 – 253.574), preeclampsia with asphyxia p-value 0.00 (OR 52.00, 95% CI 19.7 – 137.24), gestational age with infant weight at birth p-value 0.00 (OR 16.142, 95% CI 4.89 – 53.27), the Birth weight of infants with asphyxia p-value 0.00 (OR 0.329, 95% CI 0.235 – 0.459), and gestational age with asphyxia p-value 0.00 (OR 1.697, 95% CI 1.50 – 1.91).

Conclusion: Newborns can suffer from nutritional deficiencies and asphyxia due to inadequate uterine-placental blood flow and impaired uteroplacental perfusion, vasospasm, and disruption of spiral arteries while in the womb.

Keywords: Asphyxia, newborn, preeclampsia, pregnancy.

INTRODUCTION

Preeclampsia is a complication in pregnancy and is one of the causes of maternal and fetal death.^{1,2} Maternal mortality according to the Tenth Revision of The International Classification of Disease (ICD-10) is a woman's death that occurs during pregnancy, or within 42 days after the end of pregnancy.³ In developing countries, the incidence of pre-eclampsia and eclampsia ranges from 1:100 to 1:1700.⁴ The pre-eclampsia can be called a public health problem if the Case Fatality Rate (CFR) of pre-eclampsia/eclampsia reaches 1.4 – 1.8%.^{1,5} From developed countries, the maternal mortality rate is reported to be 0 – 1.8%. The perinatal mortality rate due to eclampsia in the United States and Great Britain ranges from 5.6% to 11.8%, while in developing countries the maternal mortality rate reaches 14%.⁶ Preeclampsia complications that occur in the fetus are nutritional deficiencies due to inadequate uterine-placental blood flow and can lead to premature birth, LBW, or stillbirth.^{7,8}

LBW can be an important marker for the health and nutrition of mothers and fetuses.⁹ Globally, the incidence of LBW is estimated to be around 15% of all births/year.¹⁰ Decreased uteroplacental perfusion can significantly increase the risk of intrauterine growth

restriction, preterm birth, and other neonatal comorbidities (such as respiratory distress syndrome, and intraventricular hemorrhage).^{1,10} Pre-eclampsia can also cause asphyxia in newborns which occurs due to impaired uteroplacental perfusion due to vasospasm and damage to the spiral arteries during pregnancy.^{3,11} This results in hypovolemia, vasospasm, decreased uteroplacental perfusion, and damage to the endothelial cells of the placental blood vessels.^{9,12} This study aims to analyze the risk factors for preeclampsia with birth weight and the incidence of asphyxia.

METHODOLOGY

This study used a case-control (retrospective) design with data sources of medical records in the period 2019 – 2020 for mothers who gave birth with a diagnosis of pre-eclampsia at a referral hospital in East Java, Indonesia. The variables of this study were pre-eclampsia status, baby weight, and asphyxia status at birth. The sample was 255 women who gave birth and were diagnosed with pre-eclampsia with the inclusion criteria for live births, while the exclusion criteria were pregnant women with a history of previous hypertension.

The baby weight indicator was taken from medical

records of mothers. If the birth weight was less than 2500 grams it was said to be LBW and if > 2500 was said to be normal weight. For asphyxia, if Apgar score was 1 – 3 it was labeled as severe asphyxia, 4 – 6 moderate asphyxia, and 7 – 10 mild asphyxia. The diagnosis of preeclampsia was established using blood pressure indicators; if the systolic pressure was more than 140 mmHg and the diastolic more than 80 mmHg, then pre-eclampsia was classified as mild, whereas if the systolic pressure was more than 160 mmHg with a diastolic pressure of 100 mmHg, then preeclampsia was labeled severe. Another assessment uses protein urine, as well as a history of pre-eclampsia before being referred to the hospital.

Statistical Analysis: Analysis of data was carried out using SPSS version 21 with a chi-square analysis. P-values < 0.05 were considered statistically significant and the rough/adjusted odds ratio was calculated.

RESULTS

Most of the mothers had mild pre-eclampsia 68.9%, normal baby weight (2500 – 3500 gr) 64.7%, and 70% gestational age 38 – 40 weeks (Table 1). The relationship between pre-eclampsia status and newborn body weight has a p-value of 0.00, (OR 84.00 95%, CI 27.91 - 253.574) (Table 2).

Pre-eclampsia with asphyxia p-value was 0.00 (OR 52.00, 95% CI 19.7-137.24) (Table 3), gestational age with infant weight at birth p-value 0.00 (OR 16.142, 95% CI 4.89-53.27) (Table 4), Birth weight of infants with asphyxia had p-value 0.00 (OR 0.329, 95% CI 0.235 – 0.459), and age of gestation with asphyxia had p-value 0.00 (OR 1.697, 95% CI 1.50 – 1.91). There was a relationship between preeclampsia status and the incidence of neonatal asphyxia p-value 0.00 (OR 52.00, 95% CI 19.7 – 137.24), mothers with severe preeclampsia had a 52 times risk of experiencing asphyxia.

Table 1: Age of pregnant women, gestational age, mother's education and other factors.

Characteristic	Category	n (%)
Age of pregnant women	Less than 20 years	78 (30,6)
	20 – 30 years	103(40,4)
	Over 30 years	74 (29)
Gestational age	Less than 36 weeks	76 (30)
	36 – 40 weeks (at term)	179 (70)
Mother's education	Elementary school	52 (17,7)
	Middle school	134 (45,7)
	High school	59 (20,1)
	College	10 (3,4)
Mother's job	Housewife	101 (34,5)
	Laborer	9 (3,1)
	self-employed,	141 (48,1)
	Civil servants	4 (1,4)
Pre-eclampsia type	Mild preeclampsia	204 (69,8)
	Severe preeclampsia	51 (17,4)
ANC compliance	Obey	147 (59,2)
	Disobey	108 (36,9)
Baby weight	Low	90 (35,4)
	Normal	165 (64,7)
Neonatus asphyxia status	Severe	70 (23,9)
	Moderate	185 (72,5)
Maternal weight during pregnancy	Low	8 (3,1)
	Normal	202 (79,2)
	Overweight	45 (17,6)

Table 2: Relationship between newborn weight and pre-eclampsia.

Variable	Pre-eclampsia Type		Total	OR (95% CI)	p value
	Preeclampsia Severe, Mild	Preeclampsia Mild			
Low birth weight	47	50	47	84,00 27,91 – 253,574	0,00
Normal birth weight	4	204	208		
Total	51	204	255		

Table 3: Association of neonatal asphyxia with preeclampsia.

Variable	Preeclampsia Type		Total	OR (95% CI)	p-value
	Preeclampsia Severe	Preeclampsia Mild			
Severe asphyxia	51	19	70	52,00 19,7-137,24	0,00
Moderate asphyxia	0	108	208		
Total	51	204	255		

Table 4: Relationship between gestational age and newborn body weight.

Variable	Newborn Weight		Total	OR (95% CI)	p-value
	Low	Normal			
Gestational age term (37-40 week)	3	59	63	16,142 4.89 – 53,27	0,00
Gestational age pre term (less than 36 weeks)	87	106	193		
Total	90	165	255		

DISCUSSION

Pre-eclampsia (PE) is associated with 3 major risks to the fetus which are perinatal death, intrauterine growth restriction (IUGR), and preterm birth. Perinatal mortality is increased in infants with IUGR or asphyxia.^{13,14} Premature infants with IUGR are at high risk of prolonged respiratory insufficiency.¹⁵ During pregnancy, the increased demand for oxygen causes an increase in the rate of free radical production.^{16,17} Oxidative stress is implicated in preeclampsia and IUGR.^{13,18} The level of oxidative stress and decreased antioxidant capacity may contribute to the incidence of perinatal asphyxia.¹⁵

In preeclampsia, there is a decrease in placental blood perfusion as well as vasospasm and endothelial injury.^{4,19} A placenta that experiences ischemia and hypoxia produces free radicals in the form of reactive hydroxyl radicals and lipid peroxides which circulate in the bloodstream and can damage the cell membrane, nucleus, and endothelial cell proteins which result in endothelial dysfunction.¹⁸

The results of this study indicate that there is a relationship between pre-eclampsia status with the incidence of LBW and asphyxia. Babies born with LBW and asphyxia are babies born to severe preeclampsia mothers with a gestational age of fewer than 38 weeks.²⁰ Pre-eclampsia causes mothers to give birth earlier than their gestational age. Infants born prematurely are smaller and have a higher incidence of hyaline membrane disease, patent ductus arteriosus, pulmonary air leakage, and hypotension.²¹

Efforts have been made to prevent preeclampsia during pregnancy including prophylactic aspirin therapy, which

is effective in reducing the risk of preeclampsia if given before 16 weeks of gestation.¹⁴ Other efforts that can be made to prevent preeclampsia are nutritional supplements and diet and lifestyle interventions. Vitamin D deficiency can increase the risk of preeclampsia.^{22,23} Another therapy that has been given to mothers with pre-eclampsia is the administration of MgSO₄.⁶

The only definitive treatment for preeclampsia is pregnancy termination although babies born with LBW have an increased incidence of asphyxia.²⁴ Optimal delivery time requires a careful balance of risks to both the mother and the fetus.¹⁴ If the condition of the fetus worsens, delaying labor can put the risk of impairing brain development due to prolonged fetal hypoxia, but early delivery also has risks associated with prematurity.¹⁴

CONCLUSION

Pre-eclampsia increases the incidence of preterm labor and asphyxia in newborns. There is a relationship between pre-eclampsia with LBW and asphyxia. The primary prevention program needs to be further improved through early detection of pregnant women at basic level services so that preeclampsia treatment can be carried out earlier.

Author Contributions:

Conception and design: Awatiful Azza.
 Collection and assembly of data: Awatiful Azza.
 Analysis and interpretation of data: Awatiful Azza.
 Drafting of the article: Esti Yunitasari, Cipto Susilo.
 Critical revision of article for important intellectual content: Esti Yunitasari.
 Statistical expertise: Awatiful Azza.
 Final approval and guarantor of the article: Mira Triharini.
Corresponding author email: Awatiful Azza: awatiful.azza-2020@fkip.unair.ac.id
Conflict of Interest: None declared.
 Rec. Date: Mar 4, 2022 Revision Rec. Date: May 17, 2022 Accept Date: July 2, 2022.

REFERENCES

1. Ministry of Health. Diagnosis and Treatment of Hypertension and Pre-eclampsia in Pregnancy in New Zealand: A clinical practice guideline 2018:136.
2. Guideline CP. the Diagnosis and Management of Pre-Eclampsia and Eclampsia 2013;3:1–24.
3. Xiao J, Shen F, Xue Q, Chen G, Zeng K, Stone P, et al. Is ethnicity a risk factor for developing preeclampsia? An analysis of the prevalence of preeclampsia in China. *J Hum Hypertens* 2014;28:694–8.
4. Holanda Moura BM, Lopes LM, Murthi P, Silva Costa FD. Prevention of preeclampsia. *J Pregnancy* Volume 2012 |Article ID 435090 | <https://doi.org/10.1155/2012/435090>
5. Park HJ, Kim SH, Jung YW, Shim SS, Kim JY, Cho YK, et al. Screening models using multiple markers for early detection of late-onset preeclampsia in low-risk pregnancy. *BMC Pregnancy Childbirth* 2014;14:1–11.
6. Rawlins B, Plotkin M, Rakotovo JP, Getachew A, Vaz M, Ricca J, et al. Screening and management of pre-eclampsia and eclampsia in antenatal and labor and delivery services: Findings from cross-sectional observation studies in six sub-Saharan African countries. *BMC Pregnancy Childbirth* 2018;18:1–11.
7. Zakiah N, Postma MJ, Baker PN, van Asselt ADI. Preeclampsia Diagnosis and Treatment Options: A Review of Published Economic Assessments. *Pharmacoeconomics* 2015;33:1069–82.
8. Kenny LC, Thomas G, Poston L, Myers JE, Simpson NAB, McCarthy FP, et al. Prediction of preeclampsia risk in first time pregnant women: Metabolite biomarkers for a clinical test. *PLoS One* 2020;15:1–19.
9. Galaviz-Hernandez C, Sosa-Macias M, Teran E, Garcia-Ortiz JE, Lazalde-Ramos BP. Paternal determinants in preeclampsia. *Front Physiol* 2019;10:1–7.
10. Berhe AK, Ilesanmi AO, Aimakhu CO, Mulugeta A. Effect of pregnancy-induced hypertension on adverse perinatal outcomes in Tigray regional state, Ethiopia: A prospective cohort study. *BMC Pregnancy Childbirth* 2019;20:1–11.
11. Akeju DO, Vidler M, Oladapo OT, Sawchuck D, Qureshi R, Von Dadelszen P, et al. Community perceptions of pre-eclampsia and eclampsia in Ogun State, Nigeria: A qualitative study. *Reprod Health* 2016;13.
12. Karampas GA, Eleftheriades MI, Panoulis KC, Rizou MD, Haliassos AD, Metallinou DK, et al. Prediction of pre-eclampsia combining NGAL and other biochemical markers with Doppler in the first and/or second trimester of pregnancy. A pilot study. *Eur J Obstet Gynecol Reprod Biol* 2016;205:153–7.
13. El Sayed SL, Desoky MM. Effect of lifestyle alteration of pregnant women with mild preeclampsia on maternal and fetal status. *JOJ Nurse Health Care* 2019;10:1-7.
14. de Souza Rugolo LM, Bentlin MR, Trindade CE. Preeclampsia: effect on the fetus and newborn. *Neo Reviews* 2011;12:e198-206.
15. Aslam HM uhamma., Saleem S, Afzal R, Iqbal U, Saleem SM Uhamma, Shaikh MW, et al. Risk factors of birth asphyxia. *Ital J Pediatr* 2014;40:94.
16. Poon LC, Nicolaidis KH. Early prediction of preeclampsia. *Obstet Gynecol Int* 2014;2014:297397.
17. Karampas GA, Eleftheriades MI, Panoulis KC, Rizou MD, Haliassos AD, Metallinou DK, et al. European Journal of Obstetrics & Gynecology and Reproductive Biology Prediction of pre-eclampsia combining NGAL and other biochemical markers with Doppler in the first and/or second trimester of pregnancy. A pilot study. *Eur J Obstet Gynecol* 2016;205:153–7.
18. Hypertension in pregnancy. Report of the American College of Obstetricians and Gynecologists' Task Force on Hypertension in Pregnancy. *Obstet Gynecol* 2013;122:1122-1131.
19. Dalili H, Nili F, Sheikh M, Hardani AK, Shariat M, Nayeri F. Comparison of the four proposed Apgar scoring systems in the assessment of birth asphyxia and adverse early neurologic outcomes. *PLoS One* 2015;10:1–9.
20. Wagata M, Ishiguro M, Obara T, Nagai M, Mizuno S, Nakaya N, et al. Low birth weight and abnormal pre-pregnancy body mass index were at higher risk for hypertensive disorders of pregnancy. *Pregnancy Hypertens* 2020;22:119–25.
21. Powe CE, Levine RJ, Karumanchi SA. Preeclampsia, a disease of the maternal endothelium: the role of antiangiogenic factors and implications for later cardiovascular disease. *Circulation* 2011;123:2856-69.
22. Abedi P, Mohaghegh Z, Afshari P, Latifi M. The relationship of serum vitamin D with pre-eclampsia in the Iranian women. *Matern Child Nutr* 2014;10:206–12.
23. Fogacci S, Fogacci F, Banach M, Michos ED, Hernandez A V., Lip GYH, et al. Vitamin D supplementation, and incident preeclampsia: A systematic review and meta-analysis of randomized clinical trials. *Clin Nutr* 2020;39:1742–52.
24. Park HJ, Shim SS, Cha DH. Combined screening for early detection of pre-eclampsia. *Int J Mol Sci* 2015;16:17952–74.