

SELF CARE BEHAVIOR AS AN INDICATOR

by Awatiful Azza

Submission date: 29-Feb-2020 12:17PM (UTC+0800)

Submission ID: 1266384483

File name: PSYCHOLOGICAL_ADAPTATION_IN_PATIENTS_UNDERGOING_CHEMOTHERAPY.docx (33.08K)

Word count: 2597

Character count: 14952

SELF CARE BEHAVIOR AS AN INDICATOR OF PSYCHOLOGICAL ADAPTATION IN PATIENTS UNDERGOING CHEMOTHERAPY

Awatiful Azza

Faculty of Health Sciences, University of Muhammadiyah Jember

Email: awatiful.azza@unmuhjember.ac.id

Abstract

Background: Cancer is a terminal disease and can be a threat to human well-being and health in general. One medical therapy, to prevent the development of cancer is chemotherapy. This treatment has not only positive effects but also adverse effects from sufferers.

Method: This study aims to analyze the relationship of self-care behavior with the adaptability of patients undergoing chemotherapy. The study sample was 120 patients in the chemotherapy treatment unit at the hospital in Jember-Indonesia district, with purposive sampling. Data were obtained through a modified Self Care Heart Failure In-dex (SCHFI) instrument for chemotherapy patients. Analysis of data was used in Chi-square.

The results of this study indicate there is a relationship between self-care behavior with the adaptability of patients with chemotherapy, the p-value is $0.00 < 0.005$.

Conclusion: Self-care behavior is one form of nursing service that can be applied to patients undergoing chemotherapy. The ability to engage patients independently during chemotherapy can increase patient self-esteem and will help improve patient adaptation due to the side effects of chemotherapy.

INTRODUCTION

Cancer is a terminal disease that can attack all levels of society without knowing social status, age, and gender. This disease is still a threat to human well-being and health in general(1)(2). World Health Organization (WHO) revealed an increase in the number of cancer patients each year to reach 6.25 million people, and two-thirds are from developing countries, including Indonesia(3). WHO data shows that cancer incidence increased from 12.7 million cases in 2008 to 14.1 million cases in 2012. Also, the number of deaths also increased from 7.6 million patients in 2008 to 8.2 million in 2012(3)(4). Cancer is the number 2 cause of death in the world at 13% after cardiovascular disease. Cancer incidence in 2030 is estimated to reach 26 million patients, and 17 million of them died. This condition mainly can occur in poor and developing countries that tend to be faster(2)(5).

Chemotherapy is one of the medical actions taken to treat cancer and kill cancer cells using cytostatic drugs(6). The selection of chemotherapy must be made by considering various factors, especially related to the need for a fast and significant response and quality of life(5)(7). Chemotherapy has the principle of working to inhibit and control cancer cells and kill cancer cells(2). Chemotherapy treatment can reach cancer cells that have spread to parts of the patient's body and can damage normal tissue, both acute and chronic(4). Some acute side effects, which can be caused by chemotherapy, include nausea, vomiting, alopecia, and bone marrow suppression. Whereas slow side effects, which occur vary and include pulmonary fibrosis, neuropathy, and nephropathy.

According to research conducted on 41 patients, it was found that 32% of patients had peripheral neuropathy, 14.7% felt pain in the legs, 2.4% felt numb, and 51.2% had fever after undergoing chemotherapy(7). While other studies have shown that the effects of chemotherapy cause 71.4% of chemotherapy patients complain of fatigue, and 85.7% complain of

experiencing sleep disorders(1)(6). Side effects of chemotherapy vary depending on the modification of chemotherapy drugs given. According to the National Cancer Institute, side effects that can occur due to chemotherapy using anthracycline (adriamycin/doxorubicin) chemotherapy drugs are nausea, vomiting, diarrhea, stomatitis, alopecia, susceptible to infection, thrombocytopenia, neuropathy and myalgia (8)(1). Data show that patients with chemotherapy frequency three times, with radiation frequency 12 times have energy intake, poor protein. The frequency of chemotherapy can affect nutrient intake due to side effects resulting from chemoradiation in the form of nausea, vomiting and diarrhea which can reduce the intake of nutrients for patients(9)(10). In addition to physiological effects, chemotherapy also impacts psychological changes, including helplessness, anxiety, shame, self-esteem, stress, and anger(5)(2). The psychological impact experienced by each person varies, depending on the severity (stage), type of treatment and characteristics of each patient(6)(10). About 30.0% of people living with cancer experience adjustment problems and 20.0% are diagnosed with depression. Patients with chemotherapy must be able to adapt to the treatment so that patients can receive all side effects, both physical and psychological. This condition certainly requires the support of all parties from family, friends and service providers. Self-care behavior is one form of nurse action that can be done by allowing patients to understand their condition and be directly involved in self-care during and after chemotherapy so that patients are able to improve their adaptation(11)(12). Social support helps a person live life and is needed to maintain physical and emotional well-being(13)(14). This study aims to analyze the relationship of self-care behavior with the adaptability of patients undergoing chemotherapy.

MATERIAL AND METHODOLOGY

This study uses correlation design with a cross-sectional approach, which aims to determine the relationship between self-care and the psychological adaptability of cancer patients undergoing chemotherapy.

The population is cancer patients undergoing chemotherapy at the chemotherapy center of a Jember-Indonesia hospital in the period of January - December 2018 totalling 150 patients. The sample in this study took 80% of the population as many as 120 samples. Inclusion criteria that have been set are: cancer patients in stage II or more, who are undergoing chemotherapy routinely, have health insurance. And the exclusion criteria in this study were patients undergoing chemotherapy in fragile conditions. The sampling technique used was purposive sampling.

The instrument used in this study was a questionnaire with Self Care Heart Failure In-dex (SCHFI) version 6.2 ((12)) which had been modified for patients with chemotherapy. There are three categories of self-care questionnaire namely to identify Self-Care Maintenance, Self-Care Management and Self-Care Confidence in patients with chemotherapy(12). Meanwhile, to assess adaptability, the data were obtained from an adaptation questionnaire with a Likert scale compiled by researchers. Data analysis was performed using the chi-square test with a value of $\alpha = 0.05$ and $p\text{-value} \leq \alpha$.

RESULTS

The study was conducted at a chemotherapy treatment center in a hospital in Jember district on 120 samples according to inclusion criteria. Data obtained through a questionnaire consisting of the characteristics of respondents include gender, age, education, occupation, duration of chemotherapy, marital status. As for the variables of self-care and adaptation obtained using a questionnaire.

Table 1; Demographic Data

Characteristics	n	%
-----------------	---	---

Gender :		
a. Male	48	40
b. female	72	60
Age :		
a. 15-25 years (late adolescence)	8	6,6
b. 26-35 years old (early adulthood)	27	22,5
c. More than 36 years (late adulthood)	85	70,8
Education		
a. Elementary school	39	32,5
b. Secondary school	68	56,6
c. High School	13	10,83
Occupation		
a. Did not work/Jobless	42	35
b. entrepreneur	39	32,5
c. Farmers	25	20,8
d. Employee	14	11,6
Marital status		
a. Married	78	65
b. Divorced / separated	27	22,5
c. single	15	12,5

Based on demographic data, it was found that patients who were included in the sample category were 60% female, with most patients over the age of 36 years. Meanwhile 35% did not work or jobless, and 68% married.

Table 2. Chemotherapy

Characteristics	n	%
Cancer type		
a. Breast cancer	68	56,6
b. Lymph cancer	17	14,16
c. Other cancers	35	29,12
Duration of chemotherapy treatment		
a. 6 months	32	26,6
b. 1 year	77	64,16
c. More than 1 year	11	9,16
early physical complaints during chemotherapy		
a. Nauseous vomit	100	83,3
b. Sprue	8	6,6
c. Fatigue	120	100
d. Hair loss	85	70,8
e. Dry and blackened skin	94	78,3
f. diarrhea	18	15
Patient response to the effects of chemotherapy		
a. shame	74	61,6
b. depression	72	60
c. adjustment disorder	67	55,8

The results of data analysis about the disease most of the samples suffered from breast cancer as much as 56.6%. With an average chemotherapy that has been running for 1 year which is

64.16%. In the analysis of the effects of chemotherapy on the sample obtained almost all samples undergoing chemotherapy experienced fatigue

Table 3 Analysis of self care behavior in patients undergoing chemotherapy

Dimensions of self-care behavior	Self-care behavior			
	Adequate		Inadequate	
	n	%	n	%
<i>self-care confidence</i>	83	69,1	37	30,8
<i>self-care management</i>	78	65	42	35
<i>Self care maintenance</i>	96	80	24	20

Table 4 Adaptation of patients undergoing chemotherapy

Adaptation	n	%
Adaptive	78	65
Mal adaptive	42	36
TOTAL	120	100

Based on the table above, most respondents have good adaptability.

Table 5 Relationship between self care and adaptation of patients undergoing chemotherapy

VARIABLE SELF CARE behavior	ADAPTATION OF CHEMOTHERAPY PATIENTS		P Value
	Mal adaptive	Adaptive	
In adequate	35 (29,1%)	0 (0%)	0,000
adequate	7 (5,8 %)	78 (65%)	
TOTAL	42 (35%)	78 (65%)	

Based on the correlation analysis, it was found that there was a relationship between self care behavior and client's adaptability with p value 0.00

DISCUSSION

Self care is a conceptual model of a person's ability to care for themselves independently so that the ability to maintain their health and well-being is achieved. Self care according is a natural decision-making process for selecting behaviors to maintain physiological stability (self care maintenance) and response to symptoms experienced (self-care management)(6)(15). The ability of individuals to perform self care is influenced by basic conditioning factors such as; age, sex, development status, health status, socio-cultural orientation, health care system (diagnostics, modality management), family system, lifestyle, environment and availability of resources. This is consistent with the results of this study that most respondents have a fairly mature age, so they are able to take self-care decisions well(16)(17).

Self care behavior in patients undergoing chemotherapy is a step in decision making and strategies undertaken by patients in an effort to maintain life, improve health function, and

achieve complete health to minimize the decline in quality of life. (18), states that universal self-care requisites are a major part of the life each individual lives. Activities undertaken related to universal self-care requisites are shown by maintaining adequate air, water and food that are useful for metabolism and also produce energy. Universal self care requisites can also directly affect patients with chemotherapy(12)(16). Developmental self-care requisites are an effort made to support the patient's development process when experiencing discomfort due to chemotherapy. While health deviation requisites are often associated with pain conditions experienced by patients, namely how the patient's ability to feel the condition of his pain or inability to carry out normal functions(7). The ability of self care gained through experience during chemotherapy will have an impact on lifestyle changes and can directly affect the patient's adaptation ability itself ((19)(6)). Psychosocial adaptation is a process and a final stage as an individual response to environmental stimuli to improve life goals and survive, grow, reproduce, and self-actualize ((15)). Each individual is a biopsychosocial unitary system in constant interaction with the environment, which is constantly changing(7). The results of this study indicate that the majority of respondents have adaptive behavior during chemotherapy. This condition is supported by high motivation, and his desire to recover from cancer, maturity of thinking and decision making causes patients to be able to adapt to their conditions, this is supported by respondents who have a mature age(5). In addition, the length of chemotherapy can also be a factor that supports sufferers more adaptable from the side effects of chemotherapy. Another factor affecting adaptability is marital status(18). The participation of families and people around patients in providing life support for patients undergoing chemotherapy is very significant, as indicated by 65% of the sample having a married status. The results of this study indicate that there is a relationship between self care and adaptation of patients undergoing chemotherapy with a P value of 0.00. The existence of physiological changes and chronic conditions on health is very influential on changes in a person's quality of life ((2)). Adequate support becomes very important when someone cannot meet their own needs. Someone who can not meet their own needs, often because of physical limitations or insufficient resources to overcome them. Appropriate and adequate social support can improve quality of life in patients(20)(6).

CONCLUSION

Self care behavior is a form of nursing service that can be applied to patients undergoing chemotherapy. The ability to engage patients independently during chemotherapy can increase

patient self-esteem. This will certainly make patients more adaptable to the side effects of chemotherapy.

Ethical Permit: This research has been passed through the Ethics Commission of the Faculty of Health Sciences, Muhammadiyah University Jember.

Funding Source: This study is a self-funded Research project.

Conflict of Interest: None

REFERENCE

1. American Cancer Society. Cancer Treatment & Survivorship Facts & Figures 2019-2021. Am Cancer Soc [Internet]. 2019;1-48. Available from: <https://www.cancer.org/research/cancer-facts-statistics/survivor-facts-figures.html>
2. Janelsins MC, Kohli S, Mohile SG, Usuki K, Ahles TA, Morrow GR. An update on cancer- and chemotherapy-related cognitive dysfunction: Current status. *Semin Oncol*. 2011;38(3):431-8.
3. World Health Organization. WHO Recommendations for Prevention and Treatment of Pre-Eclampsia and Eclampsia. WHO Recommendations for Prevention and Treatment of Pre-Eclampsia and Eclampsia. 2011.
4. WHO. Cancer Facts & Figures. 2016;
5. DeVita VT, Chu E. A history of cancer chemotherapy. *Cancer Res*. 2008;68(21):8643-53.
6. Periasamy U, Mohd Sidik S, Rampal L, Fadhilah SI, Akhtari-Zavare M, Mahmud R. Effect of chemotherapy counseling by pharmacists on quality of life and psychological outcomes of oncology patients in Malaysia: A randomized control trial. *Health Qual Life Outcomes*. 2017;15(1):1-10.
7. Aslam MS, Naveed S, Ahmed A, Abbas Z, Gull I, Athar MA. Side Effects of Chemotherapy in Cancer Patients and Evaluation of Patients Opinion about Starvation Based Differential Chemotherapy. *J Cancer Ther*. 2014;05(08):817-22.
8. Azza A, Susilo C, Efendi F. Supportive group therapy as a prediction of psychological adaptation of breast cancer patients undergoing chemotherapy. *Indian J Public Heal Res Dev*. 2018;9(10):441-5.
9. Cardoso F, Bedard PL, Winer EP, Pagani O, Senkus-Konefka E, Fallowfield LJ, et al. International guidelines for management of metastatic breast cancer: Combination vs sequential single-agent chemotherapy. *J Natl Cancer Inst*. 2009;101(17):1174-81.
10. Setiawan SD. The Effect Of Chemotherapy In Cancer Patient To. 2015;4:94-9.

11. Awatiful Azza I CS. The Learning Model Of Reproductive Health Through A Peer Group With A Culture Of Early Marriage In The Traditional Boarding School, Jember. In: Mohammad Afandi, editor. Bandung: Sekolah Tinggi Ilmu Kesehatan "Aisyiyah Bandung; 2014. p. 15–40. Available from: repository.stikes-aisyiyahbandung.ac.id/file.php?file=dosen...2014
12. Riegel B, Lee CS, Dickson VV, Carlson B. An update on the self-care of heart failure index. *J Cardiovasc Nurs.* 2009;24(6):485–97.
13. Classen C, Butler LD, Koopman C, Miller E, Dimiceli S, Giese-davis J, et al. Supportive-Expressive Group Therapy and Distress in Patients With Metastatic Breast Cancer. 2001;58(May).
14. Classen CC, Kraemer HC, Blasey C, Giese-davis J, Palesh OG, Atkinson A, et al. Supportive expressive group therapy for primary breast cancer patients: a randomized prospective multicenter trial. 2011;17(5):438–47.
15. Kusananto K, Sari NPWP, Harmayetty H, Efendi F, Gunawan J. Self-care model application to improve self-care agency, self-care activities, and quality of life in patients with systemic lupus erythematosus. *J Taibah Univ Med Sci [Internet].* 2018;13(5):472–8. Available from: <https://doi.org/10.1016/j.jtumed.2018.07.002>
16. Wahyuni A, Kurnia OS. Hubungan Self Care dan Motivasi dengan Kualitas Hidup Pasien Gagal Jantung Self-Care , Motivation , and Quality of Life among Patients with Heart Failure. *Stikes Fort Kock Bukittingig.* 2014;2(Vol. 2): 108–16.
17. Sandhi AK, Azza A, Komarudin. Hubungan Perubahan Citra Tubuh Dengan Kemampuan Adaptasi Psikososial Pasien Kanker Dengan Kemoterapi Di RS Tingkat III Baladhika Husada Jember. 2015;13.
18. McClelland HM. *Nursing theory: utilization & application (2nd edition).* Vol. 10, Accident and Emergency Nursing. 2002. 113 p.
19. Torre LA, Siegel RL, Ward EM, Jemal A. Global Cancer Incidence and Mortality Rates and Trends — An Update. 2016;25(January):16–28.
20. Prihatiningsih D, Sudyasih T. Perawatan Diri Pada Pasien Gagal Jantung. *J Pendidik Keperawatan Indones.* 2018;4(2).

SELF CARE BEHAVIOR AS AN INDICATOR

ORIGINALITY REPORT

6%

SIMILARITY INDEX

%

INTERNET SOURCES

6%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

Ina Nur Istiqomah, Ahmad Husein Alkaff, Mutiara Saragih, Ade Hanna Natalia, Usman Sumo Friend Tambunan. "Virtual screening of natural products as an inhibitor of DNA methyltransferase 1 enzyme for breast cancer disease", IOP Conference Series: Materials Science and Engineering, 2019

Publication

2%

2

Binu Koirala, Cheryl R. Dennison Himmelfarb, Chakra Budhathoki, Patricia M. Davidson. "Heart failure self-care, factors influencing self-care and the relationship with health-related quality of life: A cross-sectional observational study", Heliyon, 2020

Publication

2%

3

Ercole Vellone, Barbara Riegel, Antonello Cocchieri, Claudio Barbaranelli et al. "Psychometric testing of the self-care of heart failure index version 6.2", Research in Nursing & Health, 2013

Publication

1%

4

"Anticancer plants: Properties and Application",
Springer Science and Business Media LLC,
2018

Publication

1%

5

F. Cardoso. "Locally recurrent or metastatic
breast cancer: ESMO Clinical Practice
Guidelines for diagnosis, treatment and follow-
up", Annals of Oncology, 09/01/2011

Publication

1%

Exclude quotes Off

Exclude matches < 10 words

Exclude bibliography On